

Specification

Selective solid medium for the enumeration of enterobacteria, according to ISO standard 21528 and Pharmacopeial Harmonised Methods.

Presentation

10 Prepared bottles
 Bottles 250 ml
 with: 200 ± 5 ml

Packaging Details

1 box with 10 bottles 250 ml. Plastic screw inner cap.
 The use of syringes needles with a diameter greater than 0.8 mm is not recommended.

Shelf Life

12 months

Storage

2-25 °C

Composition

Composition (g/l):

Yeast extract.....	3.000
Peptone from gelatin.....	7.000
Salts bile.....	1.500
D(+) Glucose.....	10.000
Sodium chloride.....	5.000
Neutral red.....	0.030
Crystal violet.....	0.002
Agar.....	13.000

Description /Technique

Description:

This medium is a modification of the Violet Red Bile Agar and the MacConkey Agar as described by Mossel et al. The addition of glucose to the Violet Red Bile Agar enhances both the growth of the most fastidious enterobacteria and the recovery of those having suffered from adverse conditions. Mossel himself realized that by removing the lactose and keeping the glucose, the medium's efficiency remained stable.

This medium can be used as a presumptive medium for E. coli (by fluorescent reaction) if before sterilization MUG is added.

Directions for Use:

Collect, dilute and prepare samples and volumes as required according to specifications, directives, official standard regulations and/or expected results.

Melt the medium contained in the bottles in a water bath (100°C) or in a microwave oven, avoiding overheating, before pouring into Petri dishes when cooled to room temperature.

Once solidified on a flat surface, Spread the plates by streaking methodology or by spiral method. Incubate the plates right side up aerobically at 35°C± 2,0 for 24 h.

(Incubation times longer than those mentioned above, different inoculation methods or different incubation temperatures may be required depending on the sample, on the specifications,...

This medium can be inoculated directly or after any enrichment broth)

after incubation, enumerate all the reddish-violet colonies that have appeared onto the surface of the agar, with a red-violet halo due to bile salts precipitation.

Presumptive isolation of E.coli or coliforms must be confirmed by further microbiological and biochemical tests.

calculate total microbial count per ml of sample by multiplying the average number of colonies per plate by the inverse dilution factor if streaked a diluted sample. Report results as Colony Forming Unit (CFU's) per ml or g along with incubation time and temperature, that enables to differentiate total coliforms and faecal coliforms.

Note: The solid mediums can be melted in different ways: autoclave, bath and, if the customer considers appropriate, also the microwave. Whenever the microwave option is chosen, it is necessary to take certain safety measures to avoid breaking of the containers, such as loosening the screw cap and putting the bottle or tube in a water bath in the microwave. The fusion temperature and time will depend on the shape of the container, the volume of medium and the heat source. Avoid overheating as both the heating periods.

Quality control**Physical/Chemical control**

Color : Violet-pink

pH: 7.4 ± 0.2 at 25°C

Microbiological control

Melting- pour plates- Inoculate: 10-100 CFU accord. to Eur. Pharm. & 100 ± 20 CFU; min. 50 CFU (productivity)/ 10⁴-10⁶ CFU (selectivity) acc. to ISO.
 Microbiological control according to ISO 11133:2014/A1:2018.

Analytical methodology according to ISO 11133:2014/A1:2018; A2:2020.

Aerobiosis. Incubation: 30-35 °C. Reading at 24h (E.P.) / 37 ± 1 °C. Reading at 24 h (ISO)

Note: results ATCC® 8739/6538/9027 (30-35 °C) & ATCC® 8739/25922/19433/14028 (37 °C).

Microorganism

Enterococcus faecalis ATCC® 19433, WDCM 00009

Salmonella typhimurium ATCC® 14028, WDCM 00031

Ps. aeruginosa ATCC® 9027, WDCM 00026

Staphylococcus aureus ATCC® 6538, WDCM 00032

Escherichia coli ATCC® 25922, WDCM 00013

Escherichia coli ATCC® 8739, WDCM 00012 (37°C)

Escherichia coli ATCC® 8739, WDCM 00012 (32,5°C)

Growth

Inhibited

Good (50%)- Red purple colonies - Biliar precipitate

Good (50%) -Colourless colonies

Inhibited

Good (50%)- Red purple colonies - Biliar precipitate

Good (50%)- Red purple colonies - Biliar precipitate

Good (50%)- Red purple colonies - Biliar precipitate

Sterility Control

Incubation 48 h at 30-35 °C and 48 h at 20-25 °C: NO GROWTH.

Check at 7 days after incubation in same conditions.

Bibliography

- EUROPEAN PHARMACOPOEIA 8.0 (2014) 8th ed. § 2.6.13. Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. EDQM. Council of Europe. Strasbourg.
- ISO Norma 21528-1: 2004. Microbiology of food and animal feeding stuffs - Horizontal methods for the detection and enumeration of Enterobacteriaceae - Part 1: Detection and enumeration by MPN technique with pre-enrichment.
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- MOSSEL, D.A.A. (1985) Media for Enterobacteriaceae. Int. J. Food Microbiol. 2:27-35.
- MOSSEL, D.A.A., H. MENDERINK & H.H. SCHOLTS (1962) Use a Modified MacConkey Agar Medium for the selective growth and enumeration of all Enterobacteriaceae. J. Bact. 84:381.
- MOSSEL, D.A.A., M. VISER & A.M.R. CORNELISSEN (1963) The examination of foods for Enterobacteriaceae using a test of the type generally adopted for the detection of salmonellae. J. Appl. Bact. 26:444-452.
- MOSSEL, D.A.A. & M.A. RATTI (1970) Rapid detection of sub-lethally impaired cells of Enterobacteriaceae in dried foods. Appl. Microbiol. 20:273-275.
- PASCUAL ANDERSON, M^a R. (1992) Microbiología Alimentaria. Díaz de Santos, S.A. Madrid.
- USP 33 - NF 28 (2011) <62> Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. USP Corp. Inc. Rockville. MD. USA.